arXiv:1608.02760v1 [math.GR] 9 Aug 2016

ON SUPER INTEGRAL GROUPS

JUTIREKHA DUTTA AND RAJAT KANTI NATH*

Abstract. A finite non-abelian group G is called super integral if the spectrum, Laplacian spectrum and signless Laplacian spectrum of its commuting graph contain only integers. In this paper, we first compute various spectra of several families of finite non-abelian groups and conclude that those groups are super integral. As an application of our results we obtain some positive integers n such that n-centralizer groups are super integral. We also obtain some positive rational numbers r such that G is super integral if it has commutativity degree r. In the last section, we show that G is super integral if G is not isomorphic to S_4 and its commuting graph is planar. We conclude the paper showing that G is super integral if its commuting graph is toroidal.

1. Introduction

Let $A(\mathcal{G})$ and $D(\mathcal{G})$ denote the adjacency matrix and degree matrix of a graph $\mathcal{G}$ respectively. Then the Laplacian matrix and signless Laplacian matrix of $\mathcal{G}$ are given by $L(\mathcal{G}) = D(\mathcal{G}) - A(\mathcal{G})$ and $Q(\mathcal{G}) = D(\mathcal{G}) + A(\mathcal{G})$ respectively. We write Spec($\mathcal{G}$), L-Spec($\mathcal{G}$) and Q-Spec($\mathcal{G}$) to denote the spectrum, Laplacian spectrum and Signless Laplacian spectrum of $\mathcal{G}$. Also, Spec($\mathcal{G}$) $= \{\alpha_1^{a_1}, \alpha_2^{a_2}, \ldots, \alpha_l^{a_l}\}$, L-Spec($\mathcal{G}$) $= \{\beta_1^{b_1}, \beta_2^{b_2}, \ldots, \beta_m^{b_m}\}$ and Q-Spec($\mathcal{G}$) $= \{\gamma_1^{c_1}, \gamma_2^{c_2}, \ldots, \gamma_n^{c_n}\}$ where $\alpha_1, \alpha_2, \ldots, \alpha_n$ are the eigenvalues of $A(\mathcal{G})$ with multiplicities $a_1, a_2, \ldots, a_l$; $\beta_1, \beta_2, \ldots, \beta_m$ are the eigenvalues of $L(\mathcal{G})$ with multiplicities $b_1, b_2, \ldots, b_m$ and $\gamma_1, \gamma_2, \ldots, \gamma_n$ are the eigenvalues of $Q(\mathcal{G})$ with multiplicities $c_1, c_2, \ldots, c_n$ respectively.

A graph $\mathcal{G}$ is called integral or L-integral or Q-integral according as Spec($\mathcal{G}$) or L-Spec($\mathcal{G}$) or Q-Spec($\mathcal{G}$) contains only integers. The notion of integral graph was introduced by Harary and Schwenk [20] in the year 1974. A very impressive survey on integral graphs can be found in [8]. Ahmadi et al. noted that integral graphs have some interests for designing the network topology of perfect state transfer networks, see [5] and the references there in. L-integral graphs are also studied extensively over the years while Q-integral graphs are not studied much. One may conf. [3, 11, 22, 24, 32] and some of the references in [22] for several interesting results of these graphs.

Let G be a finite non-abelian group with center $Z(G)$. The commuting graph of G, denoted by Γ_G, is a simple undirected graph whose vertex set is $G \setminus Z(G)$, and two distinct vertices x and y are adjacent if and only if $xy = yx$. Various aspects of commuting graphs of different finite groups can be found in [6, 21, 25, 29]. A finite non-abelian group G is called integral or L-integral or Q-integral according as Γ_G is integral or L-integral or Q-integral. Then one may ask the following questions.

Question 1. Which finite non-abelian groups are integral?
Question 2. Which finite non-abelian groups are L-integral?
Question 3. Which finite non-abelian groups are Q-integral?

A finite non-abelian group G is called super integral if Γ_G is integral, L-integral and Q-integral. Therefore, in the line of above questions, we can also ask the following question.

Question 4. Which finite non-abelian groups are super integral?

In [15, 28], the authors have studied the spectrum of Γ_G for several families of finite non-abelian groups and determined several finite non-abelian integral groups. In this paper, we study the Laplacian spectrum and signless Laplacian spectrum of commuting graphs of those groups, viz., finite groups whose central quotient is isomorphic to $Sz(2)$ (the Suzuki group of order 20) or $\mathbb{Z}_p \times \mathbb{Z}_p$ (for

2010 *Mathematics Subject Classification.* 20D99, 05C50, 15A18, 05C25.
Key words and phrases. commuting graph, spectrum, integral graph, finite group.
*Corresponding author.

any pripe p) or D_{2m} (the dihedral group of order $2m$), quasidihedral groups, generalized quaternion groups, general linear groups, some projective special linear groups, the groups constructed by Hanaki in [19] etc. In a separate paper [16] we have studied various energies of the commuting graphs of these groups. Our computations in Section 2 reveal that all the above mentioned groups are super integral. Further, we shall show that all finite AC-groups are super integral. The rest part of this paper is devoted to some applications of the results obtained in Section 2.

Let x be an element of G. Then centralizer of x in G denoted by $C_G(x)$ is the set given by $\{y \in G : xy = yx\}$. Let $\mathrm{Cent}(G) = \{C_G(x) : x \in G\}$. A group G is called an n-centralizer group if $|\mathrm{Cent}(G)| = n$. The study of n-centralizer groups was initiated by Belcastro and Sherman [9] in the year 1994. The readers may conf. [14] for various results on n-centralizer groups. As an application of our results, in Section 3, we determine some positive integers n such that G is super integral if $|\mathrm{Cent}(G)| = n$.

The commutativity degree of G denoted by $\Pr(G)$ is the probability that a randomly chosen pair of elements of G commute. Clearly, $\Pr(G) = 1$ if and only if G is abelian. For a non-abelian group, it was shown in [18] that $\Pr(G) \leq 5/8$. Since then many mathematicians have studied this notion. The readers may conf. [10, 26] for various results on $\Pr(G)$. A survey of recent works on $\Pr(G)$ can be found in [12]. Using our results, in Section 4, we shall show that G is super integral if $\Pr(G) \in \{\frac{5}{14}, \frac{2}{5}, \frac{11}{27}, \frac{1}{2}, \frac{5}{8}\}$.

Recall that genus of a graph is the smallest non-negative integer n such that the graph can be embedded on the surface obtained by attaching n handles to a sphere. A graph is said to be planar or toroidal if the genus of the graph is zero or one respectively. It is worth mentioning that Afkhami et al. [4] and Das et al. [13] have classified all finite non-abelian groups whose commuting graphs are planar or toroidal recently. In the last section, we shall show that a finite non-abelian group G is super integral if it is not isomorphic to S_4, the symmetric group of degree 4, and the commuting graph of G is planar. We also show that a finite non-abelian group is super integral if its commuting graph is toroidal.

2. Some Computations

It is well-known that $\text{L-Spec}(K_n) = \{0^1, n^{n-1}\}$ and $\text{Q-Spec}(K_n) = \{(2n-2)^1, (n-2)^{n-1}\}$ where K_n denotes the complete graph on n vertices. Further, we have the following theorem.

Theorem 2.1. *If $\mathcal{G} = l_1K_{m_1} \sqcup l_2K_{m_2} \sqcup \cdots \sqcup l_kK_{m_k}$, where $l_iK_{m_i}$ denotes the disjoint union of l_i copies of K_{m_i} for $1 \leq i \leq k$, then*

$$\text{L-Spec}(\mathcal{G}) = \left\{0^{\sum_{i=1}^{k} l_i}, m_1^{l_1(m_1-1)}, m_2^{l_2(m_2-1)}, \ldots, m_k^{l_k(m_k-1)}\right\}$$

and

$$\begin{aligned}\text{Q-Spec}(\mathcal{G}) = \{&(2m_1-2)^{l_1}, (m_1-2)^{l_1(m_1-1)}, (2m_2-2)^{l_2}, (m_2-2)^{l_2(m_2-1)},\\ &\ldots, (2m_k-2)^{l_k}, (m_k-2)^{l_k(m_k-1)}\}.\end{aligned}$$

Now we compute the Laplacian spectrum and signless Laplacian spectrum of the commuting graphs of some families of finite non-abelian groups. We begin with some families of groups whose central factors are some well-known groups.

Theorem 2.2. *Let G be a finite group and $\frac{G}{Z(G)} \cong Sz(2)$, where $Sz(2)$ is the Suzuki group presented by $\langle a, b : a^5 = b^4 = 1, b^{-1}ab = a^2\rangle$. Then*

$$\begin{aligned}\text{L-Spec}(\Gamma_G) =&\{0^6, (4|Z(G)|)^{4|Z(G)|-1}, (3|Z(G)|)^{15|Z(G)|-5}\} \quad \textit{and}\\ \text{Q-Spec}(\Gamma_G) =&\{(8|Z(G)|-2)^1, (4|Z(G)|-2)^{4|Z(G)|-1}, (6|Z(G)|-2)^5, (3|Z(G)|-2)^{15|Z(G)|-5}\}.\end{aligned}$$

Proof. It was shown in [15, Theorem 2] that $\Gamma_G = K_{4|Z(G)|} \sqcup 5K_{3|Z(G)|}$. Therefore, by Theorem 2.1, the result follows. □

Theorem 2.3. *Let G be a finite group such that $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$, where p is a prime integer. Then*

$$\text{L-Spec}(\Gamma_G) = \{0^{p+1}, ((p-1)|Z(G)|)^{(p^2-1)|Z(G)|-p-1}\} \text{ and}$$
$$\text{Q-Spec}(\Gamma_G) = \{(2(p-1)|Z(G)|-2)^{p+1}, ((p-1)|Z(G)|-2)^{(p^2-1)|Z(G)|-p-1}\}.$$

Proof. It was shown in [28, Theorem 2.1] that $\Gamma_G = (p+1)K_{(p-1)|Z(G)|}$. Hence the result follows from Theorem 2.1. □

As a corollary we have the following result.

Corollary 2.4. *Let G be a non-abelian group of order p^3, for any prime p, then*

$$\text{L-Spec}(\Gamma_G) = \{0^{p+1}, (p^2-p)^{(p^3-2p-1}\} \text{ and}$$
$$\text{Q-Spec}(\Gamma_G) = \{(2p^2-2p-2)^{p+1}, (p^2-p-2)^{p^3-2p-1}\}.$$

Proof. Note that $|Z(G)| = p$ and $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$. Hence the result follows from Theorem 2.3. □

Theorem 2.5. *Let G be a finite group such that $\frac{G}{Z(G)} \cong D_{2m}$, for $m \geq 2$. Then*

$$\text{L-Spec}(\Gamma_G) = \{0^{m+1}, ((m-1)|Z(G)|)^{(m-1)|Z(G)|-1}, (|Z(G)|)^{m(|Z(G)|-1)}\} \text{ and}$$
$$\text{Q-Spec}(\Gamma_G) = \{(2(m-1)|Z(G)|-2)^1, ((m-1)|Z(G)|-2)^{(m-1)|Z(G)|-1},$$
$$(2|Z(G)|-2)^m, (|Z(G)|-2)^{m(|Z(G)|-1)}\}.$$

Proof. It was shown in [28, Theorem 2.5] that $\Gamma_G = K_{(m-1)|Z(G)|} \sqcup mK_{|Z(G)|}$. Hence the result follows from Theorem 2.1. □

Using Theorem 2.5, we now compute the Laplacian and signless Laplacian spectrum of the commuting graphs of the groups M_{2mn}, D_{2m} and Q_{4n} respectively.

Corollary 2.6. *Let $M_{2mn} = \langle a, b : a^m = b^{2n} = 1, bab^{-1} = a^{-1}\rangle$ be a metacyclic group, where $m > 2$. Then*

$$\text{L-Spec}(\Gamma_{M_{2mn}}) = \begin{cases} \{0^{m+1}, (mn-n)^{mn-n-1}, n^{mn-m}\} & \text{if } m \text{ is odd} \\ \{0^{\frac{m}{2}+1}, (mn-2n)^{mn-2n-1}, (2n)^{mn-\frac{m}{2}}\} & \text{if } m \text{ is even} \end{cases}$$

and

$$\text{Q-Spec}(\Gamma_{M_{2mn}}) = \begin{cases} \{(2mn-2n-2)^1, (mn-n-2)^{mn-n-1}, (2n-2)^m, (n-2)^{mn-m}\} \text{ if } m \text{ is odd} \\ \{(2mn-4n-2)^1, (mn-2n-2)^{mn-2n-1}, (4n-2)^{\frac{m}{2}}, \\ \qquad (2n-2)^{mn-\frac{m}{2}}\} \text{ if } m \text{ is even.} \end{cases}$$

Proof. Observe that $Z(M_{2mn}) = \langle b^2\rangle$ or $\langle b^2\rangle \cup a^{\frac{m}{2}}\langle b^2\rangle$ according as m is odd or even. Also, it is easy to see that $\frac{M_{2mn}}{Z(M_{2mn})} \cong D_{2m}$ or D_m according as m is odd or even. Hence, the result follows from Theorem 2.5. □

As a corollary to the above result we have the following results.

Corollary 2.7. *Let $D_{2m} = \langle a, b : a^m = b^2 = 1, bab^{-1} = a^{-1}\rangle$ be the dihedral group of order $2m$, where $m > 2$. Then*

$$\text{L-Spec}(\Gamma_{D_{2m}}) = \begin{cases} \{0^{m+1}, (m-1)^{m-2}\} & \text{if } m \text{ is odd} \\ \{0^{\frac{m}{2}+1}, (m-2)^{m-3}, 2^{\frac{m}{2}}\} & \text{if } m \text{ is even} \end{cases}$$

and

$$\text{Q-Spec}(\Gamma_{D_{2m}}) = \begin{cases} \{(2m-4)^1, (m-3)^{m-2}, (2n-2)^m, 0^m\} & \text{if } m \text{ is odd} \\ \{(2m-6)^1, (m-4)^{m-3}, 2^{\frac{m}{2}}, 0^{\frac{m}{2}}\} & \text{if } m \text{ is even.} \end{cases}$$

Corollary 2.8. *Let $Q_{4n} = \langle x, y : y^{2n} = 1, x^2 = y^n, yxy^{-1} = y^{-1}\rangle$, where $n \geq 2$, be the generalized quaternion group of order $4n$. Then*

$$\text{L-Spec}(\Gamma_{Q_{4n}}) = \{0^{n+1}, (2n-2)^{2n-3}, 2^n\} \text{ and } \text{Q-Spec}(\Gamma_{Q_{4n}}) = \{(4n-6)^1, (2n-4)^{2n-3}, 2^n, 0^n\}.$$

Proof. The result follows from Theorem 2.5 noting that $Z(Q_{4n}) = \{1, a^n\}$ and $\frac{Q_{4n}}{Z(Q_{4n})} \cong D_{2n}$. □

Now we compute the Laplacian spectrum and signless Laplacian spectrum of the commuting graphs of some well-known families of finite non-abelian groups.

Proposition 2.9. *Let G be a non-abelian group of order pq, where p and q are primes with $p \mid (q-1)$. Then*

$$\begin{aligned}\text{L-Spec}(\Gamma_G) =& \{0^{q+1}, (q-1)^{q-2}, (p-1)^{pq-2q}\} \quad \text{and}\\ \text{Q-Spec}(\Gamma_G) =& \{(2q-4)^1, (q-3)^{q-2}, (2p-4)^q, (p-3)^{pq-2q}\}.\end{aligned}$$

Proof. It was shown in [15, Lemma 3] that $\Gamma_G = K_{q-1} \sqcup qK_{p-1}$. Hence, the result follows from Theorem 2.1. □

Proposition 2.10. *Let QD_{2^n} denote the quasidihedral group $\langle a, b : a^{2^{n-1}} = b^2 = 1, bab^{-1} = a^{2^{n-2}-1}\rangle$, where $n \geq 4$. Then*

$$\begin{aligned}\text{L-Spec}(\Gamma_{QD_{2^n}}) =& \{0^{2^{n-2}+1}, (2^{n-1}-2)^{2^{n-1}-3}, 2^{2^{n-2}}\} \quad \text{and}\\ \text{Q-Spec}(\Gamma_{QD_{2^n}}) =& \{(2^n-6)^1, (2^{n-1}-4)^{2^{n-1}-3}, 2^{2^{n-2}}, 0^{2^{n-2}}\}.\end{aligned}$$

Proof. It was shown in [15, Proposition 1] that $\Gamma_{QD_{2^n}} = K_{2^{n-1}-2} \sqcup 2^{n-2}K_2$. Hence, the result follows from Theorem 2.1. □

Proposition 2.11. *The Laplacian spectrum and signless Laplacian spectrum of the commuting graph of the projective special linear group $PSL(2, 2^k)$, where $k \geq 2$, are given by*

$$\{0^{2^{2k}+2^k+1}, (2^k-1)^{2^{2k}-2^k-2}, (2^k-2)^{2^{k-1}(2^{2k}-2^{k+1}-3)}, (2^k)^{2^{k-1}(2^{2k}-2^{k+1}+1)}\} \text{ and}$$

$$\{(2^{k+1}-4)^{2^k+1}, (2^k-3)^{2^{2k}-2^k-2}, (2^{k+1}-6)^{2^{k-1}(2^k+1)}, (2^k-4)^{2^{k-1}(2^{2k}-2^{k+1}-3)}, (2^{k+1}-2)^{2^{k-1}(2^k-1)}, (2^k-2)^{2^{k-1}(2^{2k}-2^{k+1}+1)}\} \text{ respectively.}$$

Proof. It was shown in [15, Proposition 2] that

$$\Gamma_{PSL(2,2^k)} = (2^k+1)K_{2^k-1} \sqcup 2^{k-1}(2^k+1)K_{2^k-2} \sqcup 2^{k-1}(2^k-1)K_{2^k}.$$

Hence, the result follows from Theorem 2.1. □

Proposition 2.12. *The Laplacian spectrum and signless Laplacian spectrum of the commuting graph of the general linear group $GL(2, q)$, where $q = p^n > 2$ and p is a prime integer, are given by*

$$\{0^{q^2+q+1}, (q^2-3q+2)^{\frac{q(q+1)(q^2-3q+1)}{2}}, (q^2-q)^{\frac{q(q-1)(q^2-q-1)}{2}}, (q^2-2q+1)^{q(q+1)(q-2)}\} \text{ and}$$

$$\{(2q^2-6q-2)^{\frac{q(q+1)}{2}}, (q^2-3q)^{\frac{q(q+1)(q^2-3q+1)}{2}}, (2q^2-2q-2)^{\frac{q(q-1)}{2}}, (q^2-q-2)^{\frac{q(q-1)(q^2-q-1)}{2}}, (2q^2-4q)^{q+1}, (q^2+2q-1)^{q(q+1)(q-2)}\} \text{ respectively.}$$

Proof. It was shown in [15, Proposition 3] that

$$\Gamma_{GL(2,q)} = \frac{q(q+1)}{2}K_{q^2-3q+2} \sqcup \frac{q(q-1)}{2}K_{q^2-q} \sqcup (q+1)K_{q^2-2q+1}.$$

Hence, the result follows from Theorem 2.1. □

Proposition 2.13. *Let $F = GF(2^n), n \geq 2$ and ϑ be the Frobenius automorphism of F, i. e., $\vartheta(x) = x^2$ for all $x \in F$. Let $A(n, \vartheta)$ denote the group*

$$\left\{ U(a,b) = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & \vartheta(a) & 1 \end{bmatrix} : a, b \in F \right\}.$$

under matrix multiplication given by $U(a,b)U(a',b') = U(a+a', b+b'+a'\vartheta(a))$. Then

$$\text{L-Spec}(\Gamma_{A(n,\vartheta)}) = \{0^{2^n-1}, (2^n)^{2^{2n}-2^{n+1}+1}\} \text{ and}$$
$$\text{Q-Spec}(\Gamma_{A(n,\vartheta)}) = \{(2^{n+1}-2)^{2^n-1}, (2^n-2)^{2^{2n}-2^{n+1}+1}\}.$$

Proof. It was shown in [15, Proposition 4] that $\Gamma_{A(n,\vartheta)} = (2^n-1)K_{2^n}$. Hence the result follows from Theorem 2.1. □

Proposition 2.14. *Let $F = GF(p^n)$ where p is a prime. Let $A(n,p)$ denote the group*

$$\left\{ V(a,b,c) = \begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & c & 1 \end{bmatrix} : a, b, c \in F \right\}.$$

under matrix multiplication $V(a,b,c)V(a',b',c') = V(a+a', b+b'+ca', c+c')$. Then

$$\text{L-Spec}(\Gamma_{A(n,p)}) = \{0^{p^n+1}, (p^{2n}-p^n)^{p^{3n}-2p^n-1}\} \text{ and}$$
$$\text{Q-Spec}(\Gamma_{A(n,p)}) = \{(2p^{2n}-2p^n-2)^{p^n+1}, (p^{2n}-p^n-2)^{p^{3n}-2p^n-1}\}.$$

Proof. It was shown in [15, Proposition 1] that $\Gamma_{A(n,p)} = (p^n+1)K_{p^{2n}-p^n}$. Hence the result follows from Theorem 2.1. □

Our computations reveal that all the groups considered above are both L-integral and Q-integral. Also, it was shown in [15, 28] that theses groups are integral. Hence, all these groups are super integral.

A group G is called an AC-group if $C_G(x)$ is abelian for all $x \in G \setminus Z(G)$. Various aspects of AC-groups can be found in [1, 13, 30]. In [15], the authors have shown that finite AC-groups are integral. In the following two results we shall show that if G is an AC-group or G is isomorphic to $H \times A$, where H is a finite non-abelian AC-group and A is any finite abelian group then G is L-integral as well as Q-integral. Hence, G is super integral.

Theorem 2.15. *Let G be a finite non-abelian AC-group. Then*

$$\text{L-Spec}(\Gamma_G) = \{0^n, (|X_1|-|Z(G)|)^{|X_1|-|Z(G)|-1}, \ldots, (|X_n|-|Z(G)|)^{|X_n|-|Z(G)|-1}\} \text{ and}$$
$$\text{Q-Spec}(\Gamma_G) = \{(2(|X_1|-|Z(G)|)-2)^1, (|X_1|-|Z(G)|-2)^{|X_1|-|Z(G)|-1}, \ldots,$$
$$(2(|X_n|-|Z(G)|)-2)^1, (|X_n|-|Z(G)|-2)^{|X_n|-|Z(G)|-1}\}.$$

where $X_1, \ldots, X_n$ are the distinct centralizers of non-central elements of G.

Proof. By Lemma 1 in [15], we have $\Gamma_G = \bigsqcup_{i=1}^{n} K_{|X_i|-|Z(G)|}$. Therefore, the result follows from Theorem 2.1. □

Corollary 2.16. *Let $G \cong H \times A$ where H is a finite non-abelian AC-group and A is any finite abelian group. Then*

$$\text{L-Spec}(\Gamma_G) = \{0^n, (|A|(|X_1|-|Z(H)|))^{|A|(|X_1|-|Z(H)|)-1}, \ldots,$$
$$(|A|(|X_n|-|Z(H)|))^{|A|(|X_n|-|Z(H)|)-1}\} \text{ and}$$
$$\text{Q-Spec}(\Gamma_G) = \{(2|A|(|X_1|-|Z(H)|)-2)^1, (|A|(|X_1|-|Z(H)|)-2)^{|A|(|X_1|-|Z(H)|)-1}, \ldots,$$
$$(2|A|(|X_n|-|Z(H)|)-2)^1, (|A|(|X_n|-|Z(H)|)-2)^{|A|(|X_n|-|Z(H)|)-1}\}.$$

where $X_1, \ldots, X_n$ are the distinct centralizers of non-central elements of H.

Proof. It is easy to see that $Z(H \times A) = Z(H) \times A$ and $X_1 \times A, X_2 \times A, \ldots, X_n \times A$ are the distinct centralizers of non-central elements of $H \times A$. Therefore, if H is an AC-group then $H \times A$ is also an AC-group. Hence, the result follows from Theorem 2.15. □

3. n-CENTRALIZER GROUPS

For a finite group G, the set $C_G(x) = \{y \in G : xy = yx\}$ is called the centralizer of an element $x \in G$. Let $|\operatorname{Cent}(G)| = |\{C_G(x) : x \in G\}|$, that is the number of distinct centralizers in G. A group G is called an n-centralizer group if $|\operatorname{Cent}(G)| = n$. The study of these groups was initiated by Belcastro and Sherman [9] in the year 1994. The readers may conf. [14] for various results on these groups. In this section, we ask the following question.

Question 5. Can we determine all the positive integers n such that n-centralizer groups are super integral?

In the next three results we shall show that an n-centralizer group is super integral if $n = 4, 5$. Further, we shall show that a $(p+2)$-centralizer p-group is super integral for any prime p.

Theorem 3.1. *A finite* 4*-centralizer group is super integral.*

Proof. Let G be a finite 4-centralizer group. Then, by [9, Theorem 2], we have $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. Therefore, by Theorem 2.3, we have

$$\text{L-Spec}(\Gamma_G) = \{0^3, (|Z(G)|)^{3|Z(G)|-3}\} \text{ and } \text{Q-Spec}(\Gamma_G) = \{(2|Z(G)|-2)^3, (|Z(G)|-2)^{3|Z(G)|-3}\}.$$

Also, by [15, Theorem 3], we have $\text{Spec}(\Gamma_G) = \{(-1)^{3|Z(G)|-3}, (|Z(G)|-1)^3\}$. Hence, G is super integral group. □

Further, we have the following result.

Corollary 3.2. *For any prime p, a finite $(p+2)$-centralizer p-group is super integral.*

Proof. Let G be a finite $(p+2)$-centralizer p-group. Then, by [7, Lemma 2.7], we have $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$. Therefore, by Theorem 2.3, we have

$$\begin{aligned}\text{L-Spec}(\Gamma_G) =& \{0^{p+1}, ((p-1)|Z(G)|)^{(p^2-1)|Z(G)|-p-1}\} \text{ and}\\ \text{Q-Spec}(\Gamma_G) =& \{(2(p-1)|Z(G)|-2)^{p+1}, ((p-1)|Z(G)|-2)^{(p^2-1)|Z(G)|-p-1}\}.\end{aligned}$$

Also, by [28, Corollary 2.4], we have $\text{Spec}(\Gamma_G) = \{(-1)^{(p^2-1)|Z(G)|-p-1}, ((p-1)|Z(G)|-1)^{p+1}\}$. Hence, G is a super integral group. □

Theorem 3.3. *A finite* 5*-centralizer group is super integral.*

Proof. Let G be a finite 5-centralizer group. Then by [9, Theorem 4] we have $\frac{G}{Z(G)} \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ or D_6. Now, if $\frac{G}{Z(G)} \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ then by Theorem 2.3 we have $\text{L-Spec}(\Gamma_G) = \{0^4, (2|Z(G)|)^{8|Z(G)|-4}\}$ and $\text{Q-Spec}(\Gamma_G) = \{4|Z(G)|-2)^4, (2|Z(G)|-2)^{8|Z(G)|-4}\}$. If $\frac{G}{Z(G)} \cong D_6$ then by Theorem 2.5 we have $\text{L-Spec}(\Gamma_G) = \{0^4, (2|Z(G)|)^{2|Z(G)|-1}, (|Z(G)|)^{3(|Z(G)|-1)}\}$ and $\text{Q-Spec}(\Gamma_G) = \{(4|Z(G)|-2)^1, (2|Z(G)|-2)^{2|Z(G)|-1}, (2|Z(G)|-2)^3, (|Z(G)|-2)^{3(|Z(G)|-1)}\}$. Therefore, G is both L-integral and Q-integral. Also, in [28, Corollary 2.6], it was shown that a finite 5-centralizer group G is integral. Hence, G is super integral. □

We conclude this section with the following corollary.

Corollary 3.4. *Let G be a finite non-abelian group and $\{x_1, x_2, \ldots, x_r\}$ be a set of pairwise non-commuting elements of G having maximal size. Then G is super integral if $r = 3, 4$.*

Proof. By Lemma 2.4 in [2], we have that G is a 4-centralizer or a 5-centralizer group according as $r = 3$ or 4. Hence the result follows from Theorem 3.1 and Theorem 3.3. □

4. Commutativity degree and super integral group

Let G be a finite group. The commutativity degree of G is given by the ratio

$$\Pr(G) = \frac{|\{(x,y) \in G \times G : xy = yx\}|}{|G|^2}.$$

The origin of the commutativity degree of a finite group lies in a paper of Erdös and Turán (see [17]). Readers may conf. [10, 12, 26] for various results on $\Pr(G)$. In this section, we ask the following question.

Question 6. Can we determine all the positive rational numbers r such that any group G with $\Pr(G) = r$ is super integral?

The following theorems give some rational numbers r such that G is super integral if $\Pr(G) = r$.

Theorem 4.1. *If* $\Pr(G) \in \{\frac{5}{14}, \frac{2}{5}, \frac{11}{27}, \frac{1}{2}, \frac{5}{8}\}$ *then* G *is super integral.*

Proof. If $\Pr(G) \in \{\frac{5}{14}, \frac{2}{5}, \frac{11}{27}, \frac{1}{2}, \frac{5}{8}\}$ then as shown in [31, pp. 246] and [27, pp. 451], we have $\frac{G}{Z(G)}$ is isomorphic to one of the groups in $\{D_{14}, D_{10}, D_8, D_6, \mathbb{Z}_2 \times \mathbb{Z}_2\}$. If $\frac{G}{Z(G)}$ is isomorphic to D_{14}, D_{10}, D_8 or D_6 then by [28, Theorem 2.5], we have G is integral and by Theorem 2.5, we have G is both L-integral and Q-integral. Hence, in this case G is super integral. If $\frac{G}{Z(G)}$ is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$ then by [15, Theorem 3], it follows that G is integral. Also, by Theorem 2.3 we have G is both L-integral and Q-integral. Hence, G is super integral. This completes the proof. □

Theorem 4.2. *Let* G *be a finite group and* p *the smallest prime divisor of* $|G|$. *If* $\Pr(G) = \frac{p^2+p-1}{p^3}$ *then* G *is super integral.*

Proof. If $\Pr(G) = \frac{p^2+p-1}{p^3}$ then by [23, Theorem 3] we have $\frac{G}{Z(G)}$ is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$. Now, by [28, Theorem 2. 1], we have that G is integral. Again, by Theorem 2.3, it follows that G is L-integral as well as Q-integral. Hence, G is super integral. □

We conclude this section by the following result.

Theorem 4.3. *If* G *is a non-solvable group with* $\Pr(G) = \frac{1}{12}$ *then* G *is super integral.*

Proof. By Proposition 3.3.7 in [10] we have that G is isomorphic to $A_5 \times B$ for some abelian group B. Therefore G is an AC-group and hence super integral. □

5. More Applications

In this section, we show that a finite non-abelian group G not isomorphic to S_4 is super integral if its commuting graph is planar. We also show that G is super integral if its commuting graph is toroidal. We begin with the following useful lemma.

Lemma 5.1. *Let* G *be a group isomorphic to any of the following groups*

(1) $\mathbb{Z}_2 \times D_8$
(2) $\mathbb{Z}_2 \times Q_8$
(3) $M_{16} = \langle a, b : a^8 = b^2 = 1, bab = a^5 \rangle$
(4) $\mathbb{Z}_4 \rtimes \mathbb{Z}_4 = \langle a, b : a^4 = b^4 = 1, bab^{-1} = a^{-1} \rangle$
(5) $D_8 * \mathbb{Z}_4 = \langle a, b, c : a^4 = b^2 = c^2 = 1, ab = ba, ac = ca, bc = a^2cb \rangle$
(6) $SG(16,3) = \langle a, b : a^4 = b^4 = 1, ab = b^{-1}a^{-1}, ab^{-1} = ba^{-1} \rangle$.

Then L-Spec$(\Gamma_G) = \{0^3, 4^9\}$ *and* Q-Spec$(\Gamma_G) = \{6^3, 2^9\}$.

Proof. If G is isomorphic to any of the above listed groups, then $|G| = 16$ and $|Z(G)| = 4$. Therefore, $\frac{G}{Z(G)} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. Thus the result follows from Theorem 2.3. □

Now we state and proof the main results of this section.

Theorem 5.2. *Let* Γ_G *be the commuting graph of a finite non-abelian group* G. *If* G *is not isomorphic to* S_4 *and* Γ_G *is planar then* G *is super integral.*

Proof. It was shown in [15, Theorem 4] that G is an integral group if it is not isomorphic to S_4 and Γ_G is planar.

By [4, Theorem 2.2], we have that Γ_G is planar if and only if G is isomorphic to either D_6, D_8, D_{10}, $D_{12}, Q_8, Q_{12}, \mathbb{Z}_2 \times D_8, \mathbb{Z}_2 \times Q_8, M_{16}, \mathbb{Z}_4 \rtimes \mathbb{Z}_4, D_8 * \mathbb{Z}_4, SG(16,3), A_4, A_5, S_4, SL(2,3)$ or $Sz(2)$.

If $G \cong D_6, D_8, D_{10}$ or D_{12} then by Corollary 2.7, one may conclude that Γ_G is both L-integral and Q-integral. Hence, G is both L-integral Q-integral. If $G \cong Q_8$ or Q_{12} then G is both L-integral and Q-integral, by Corollary 2.8. If $G \cong \mathbb{Z}_2 \times D_8, \mathbb{Z}_2 \times Q_8, M_{16}, \mathbb{Z}_4 \rtimes \mathbb{Z}_4, D_8 * \mathbb{Z}_4$ or $SG(16,3)$ then, by Lemma 5.1, G is L-integral as well as Q-integral.

If $G \cong A_4 = \langle a, b : a^2 = b^3 = (ab)^3 = 1\rangle$ then the distinct centralizers of non-central elements of G are $C_G(a) = \{1, a, bab^2, b^2ab\}, C_G(b) = \{1, b, b^2\}$, $C_G(ab) = \{1, ab, b^2a\}, C_G(ba) = \{1, ba, ab^2\}$ and $C_G(aba) = \{1, aba, bab\}$. Note that these centralizers are abelian subgroups of G. Therefore, $\Gamma_G = K_3 \sqcup 4K_2$. Using Theorem 2.1, we have L-Spec$(\Gamma_G) = \{0^5, 3^2, 2^4\}$ and Q-Spec$(\Gamma_G) = \{4^1, 1^2, 2^4, 0^4\}$. Hence, G is both L-integral and Q-integral.

If $G \cong Sz(2)$ then by Theorem 2.2, we have L-Spec$(\Gamma_G) = \{0^6, 4^3, 3^{10}\}$ and Q-Spec$(\Gamma_G) = \{6^1, 2^3, 4^5, 1^{10}\}$. Hence, G is both L-integral and Q-integral.

If G is isomorphic to $SL(2,3)$ then it was shown in the proof of [15, Theorem 4] that $\Gamma_G = 3K_2 \sqcup 4K_4$. Therefore, by Theorem 2.1, we have L-Spec$(\Gamma_G) = \{0^7, 2^3, 4^{12}\}$ and Q-Spec$(\Gamma_G) = \{0^3, 2^{15}, 6^4\}$. Hence, G is both L-integral and Q-integral.

If $G \cong A_5$ then by Proposition 2.11, we have

$$\text{L-Spec}(\Gamma_G) = \{0^{21}, 3^{10}, 2^{10}, 4^{18}\} \text{ and } \text{Q-Spec}(\Gamma_G) = \{4^5, 1^{10}, 2^{10}, 0^{10}, 6^6, 2^{18}\},$$

noting that $PSL(2,4) \cong A_5$. Hence, G is both L-integral and Q-integral.

Finally, if $G \cong S_4$ then it can be seen that the characteristic polynomial of $L(\Gamma_G)$ is $x^5(x-1)^3(x-2)^4(x-3)^6(x-5)(x^2-8x+3)^2$ and so

$$\text{L-Spec}(\Gamma_G) = \left\{0^5, 1^3, 2^4, 3^6, 5^1(4+\sqrt{13})^2, (4-\sqrt{13})^2\right\}.$$

Also, the characteristic polynomial of $Q(\Gamma_G)$ is $x^4(x-1)^6(x-2)^4(x-3)^3(x^2-11x+20)(x^2-8x+11)^2$ and so

$$\text{Q-Spec}(\Gamma_G) = \left\{0^4, 1^6, 2^4, 3^3, 5^1(4+\sqrt{5})^2, (4-\sqrt{5})^2, \left(\frac{11+\sqrt{41}}{2}\right)^1, \left(\frac{11-\sqrt{41}}{2}\right)^1\right\}.$$

This shows that if $G \cong S_4$ then it is neither L-integral nor Q-integral. This completes the proof. □

Theorem 5.3. *Let Γ_G be the commuting graph of a finite non-abelian group G. Then G is super integral if Γ_G is toroidal.*

Proof. By Theorem 6.6 of [13], we have Γ_G is toroidal if and only if G is isomorphic to either $D_{14}, D_{16}, Q_{16}, QD_{16}, D_6 \times \mathbb{Z}_3, A_4 \times \mathbb{Z}_2$ or $\mathbb{Z}_7 \rtimes \mathbb{Z}_3$.

If $G \cong D_{14}$ or D_{16} then, by Corollary 2.7, one may conclude that G is both L-integral and Q-integral. If $G \cong Q_{16}$ then, by Corollary 2.8, G becomes both L-integral and Q-integral. If $G \cong QD_{16}$ then, by Proposition 2.10, G becomes both L-integral and Q-integral. If $G \cong \mathbb{Z}_7 \rtimes \mathbb{Z}_3$ then G is both L-integral and Q-integral, by Proposition 2.9. If G is isomorphic to $D_6 \times \mathbb{Z}_3$ or $A_4 \times \mathbb{Z}_2$ then G becomes both L-integral and Q-integral by Corollary 2.16, since D_6 and A_4 are AC-groups. Further, it was shown in [15, Theorem 5] that G is integral if Γ_G is toroidal. Hence, G is super integral if Γ_G is toroidal. □

We conclude the paper with the following result.

Proposition 5.4. *Let Γ_G be the commuting graph of a finite non-abelian group G. Then Γ_G is super integral if the complement of Γ_G is planar.*

Proof. If the complement of Γ_G is planar then, by [1, Proposition 2.3], we have G is isomorphic to either D_6, D_8 or Q_8. If $G \cong D_6$ or D_8 then, by Corollary 2.7 and [15, Proposition 6], we have that

G is super integral. If $G \cong Q_8$ then, by Corollary 2.8 and [15, Proposition 7], it follows that G is super integral. This completes the proof. □

References

[1] A. Abdollahi, S. Akbari and H. R. Maimani, Non-commuting graph of a group, *J. Algebra*, **298**, 468–492 (2006).
[2] A. Abdollahi, S. M. Jafarain and A. M. Hassanabadi, Groups with specific number of centralizers, *Houston J. Math.*, **33**(1), 43–57 (2007).
[3] N. M. M. Abreu, C. T. M. Vinagre, A. S. Bonifácioa and I. Gutman, The Laplacian energy of some Laplacian integral graph, *MATCH Commun. Math. Comput. Chem.*, **60**, 447–460 (2008)
[4] M. Afkhami, M. Farrokhi D. G. and K. Khashyarmanesh, Planar, toroidal, and projective commuting and non-commuting graphs, *Comm. Algebra*, **43**(7), 2964–2970 (2015).
[5] O. Ahmadi, N. Alon, I. F. Blake and I. E. Shparlinski, Graphs with integral spectrum, *Linear Algebra Appl.*, **430**(1), 547–552 (2009).
[6] S. Akbari, A. Mohammadian, H. Radjavi and P. Raja, On the diameters of commuting graphs, *Linear Algebra Appl.*, **418**, 161–176 (2006).
[7] A. R. Ashrafi, On finite groups with a given number of centralizers, *Algebra Colloq.*, **7**(2), 139–146 (2000).
[8] K. Balińska, D. Cvetković, Z. Radosavljević, S. Simić and D. Stevanović, A survey on integral graphs, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat.*, **13**, 42–65 (2003).
[9] S. M. Belcastro and G. J. Sherman, Counting centralizers in finite groups, *Math. Magazine*, **67**(5), 366–374 (1994).
[10] A. Castelaz, *Commutativity degree of finite groups*, M.A. thesis, Wake Forest University (2010).
[11] D. Cvetković, P. Rowlinson, S. Simić, Signless Laplacian of finite graphs, *Linear Algebra Appl.*, **423**, 155–171 (2007).
[12] A. K. Das, R. K. Nath and M. R. Pournaki, A survey on the estimation of commutativity in finite groups, *Southeast Asian Bull. Math.*, **37**(2), 161–180 (2013).
[13] A. K. Das and D. Nongsiang, On the genus of the commuting graphs of finite non-abelian groups, *Int. Electron. J. Algebra*, **19**, 91–109 (2016).
[14] J. Dutta, *A study of finite groups in terms of their centralizers*, M. Phil. thesis, North-Eastern Hill University (2010).
[15] J. Dutta and R. K. Nath, Spectrum of commuting graphs of some classes of finite groups, *Matematika* to appear.
[16] J. Dutta and R. K. Nath, Various energies of some super integral groups, preprint.
[17] P. Erdös and P. Turán, On some problems of a statistical group-theory IV, *Acta. Math. Acad. Sci. Hungar.*, **19**, 413–435 (1968).
[18] W. H. Gustafson, What is the probability that two group elements commute?, *Amer. Math. Monthly*, **80**, 1031–1034 (1973).
[19] A. Hanaki, A condition of lengths of conjugacy classes and character degree, *Osaka J. Math.*, **33**, 207–216 (1996).
[20] F. Harary and A. J. Schwenk, Which graphs have integral spectra?, *Graphs and Combin.*, Lect. Notes Math., Vol 406, Springer-Verlag, Berlin, 45–51 (1974).
[21] A. Iranmanesh and A. Jafarzadeh, Characterization of finite groups by their commuting graph, *Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis*, **23**(1), 7–13 (2007).
[22] S. Kirkland, Constructably Laplacian integral graphs, *Linear Algebra Appl.*, **423**, 3–21 (2007).
[23] D. MacHale, How commutative can a non-commutative group be?, *Math. Gaz.*, **58**, 199–202 (1974).
[24] R. Merris, Degree maximal graphs are Laplacian integral, *Linear Algebra Appl.*, **199**, 381–389 (1994).
[25] G. L. Morgan and C. W. Parker, The diameter of the commuting graph of a finite group with trivial center, *J. Algebra*, **393**(1), 41–59 (2013).
[26] R. K. Nath, *Commutativity degrees of finite groups – a survey*, M. Phil. thesis, North-Eastern Hill University (2008).
[27] R. K. Nath, Commutativity degree of a class of finite groups and consequences. *Bull. Aust. Math. Soc.*, **88**(3), 448–452 (2013).
[28] R. K. Nath and J. Dutta, Finite groups whose commuting graphs are integral, submitted for publication.
[29] C. Parker, The commuting graph of a soluble group, *Bull. London Math. Soc.*, **45**(4), 839–848 (2013).
[30] D. M. Rocke, p-groups with abelian centralizers, *Proc. London Math. Soc.*, **30**(3), 55–75 (1975).
[31] D. J. Rusin, What is the probability that two elements of a finite group commute?, *Pacific J. Math.*, **82**(1), 237–247 (1979).
[32] S. K. Simić and Z. Stanić, Q-integral graphs with edge-degrees at most five, *Discrete Math.*, **308**, 4625–4634 (2008).

Department of Mathematical Sciences, Tezpur University, Napaam-784028, Sonitpur, Assam, India.
E-mail address: jutirekhadutta@yahoo.com, rajatkantinath@yahoo.com*